

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Terms	Documents
06277693 or 09215996	2

Database: US Patents Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Refine Search: 06277693 or 09215996 Clear

Search History

Today's Date: 11/24/2000

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
JPAB	06277693 or 09215996	2	<u>L12</u>
JPAB	6277693 or 9215996	0	<u>L11</u>
JPAB	6277693 or 9215996	0	<u>L10</u>
USPT	15 not 16	50	<u>L9</u>
USPT	17 and 11	0	<u>L8</u>
USPT	((424/93.46)!.CCLS.)	87	<u>L7</u>
USPT	13 and isolat\$.clm.	51	<u>L6</u>
USPT	13 and isolat\$	101	<u>L5</u>
USPT	13 and acclimat\$	0	<u>L4</u>
USPT	12 and (9 or 10 or 11)	108	<u>L3</u>
USPT	11 and pH	108	<u>L2</u>
USPT	alkalophilus	110	<u>L1</u>

=> d hist

(FILE 'HOME' ENTERED AT 16:10:20 ON 24 NOV 2000)

FILE 'CAPLUS, MEDLINE, BIOSIS' ENTERED AT 16:10:41 ON 24 NOV 2000

L1	2534 S CLA
L2	7441 S OCTADECADIENOI?
L3	9804 S L1 OR L2
L4	1125 S L3 AND CONJUGAT?
L5	100 S L4 AND ISOMERIZ?
L6	49 S L5 AND ESTER?
L7	32 DUPLICATE REMOVE L6 (17 DUPLICATES REMOVED)

L5 ANSWER 5 OF 39 CAPLUS COPYRIGHT 2000 ACS

ACCESSION NUMBER: 1987:182615 CAPLUS

DOCUMENT NUMBER: 106:182615

TITLE: Method for preparation of chromatographically pure
phosphatidylcholine from commercial
sunflower lecithin

AUTHOR(S): Stefanov, K.; Seizova, K.

CORPORATE SOURCE: Inst. Org. Chem., Sofia, 1040, Bulg.

SOURCE: Izv. Khim. (1986), 19(4), 528-33

CODEN: IZKHDX; ISSN: 0324-0401

DOCUMENT TYPE: Journal

LANGUAGE: Bulgarian

AB Com. **sunflower** lecithin in general contains 40-50% neutral and 35-45% polar lipids, e.g., **phosphatidycholine** (PC), **phosphatidylethanolamine** (PE) and **phosphatidylinositol** (PI) in approx. equal quantities. The isolation of PC-enriched fraction was based on the acylation of PE and partly of PI at room temp. in a CHCl₃-pyridine soln. with a 5-fold excess of Ac₂O. The reaction mixt.

was

pptd. with Me₂CO and dissolved in EtOH. The soln. contg. >85% PC can be used directly as a starting material for the prepn. of liposomes or purified by column chromatog. on silica or on alumina and eluted by CHCl₃-MeOH.